

All posttests must be submitted online to AAPA.

AAPA members should complete and submit posttests on the AAPA Web site by going to [www.aapa.org](http://www.aapa.org) and searching for keyword *JAAPA post-tests*. All others may complete and submit posttests online at no charge at [www.mycme.com](http://www.mycme.com). To obtain 1 hour of AAPA Category I CME credit, PAs must receive a score of 70% or better on each test taken.

### LYME DISEASE, PAGE 22

1. The classic rash associated with Lyme disease manifests at the site of inoculation, commonly
  - a. On the back
  - b. At the waistline
  - c. In the popliteal area
  - d. In the neck region
2. The monoarticular process in stage 2 or 3 Lyme disease most commonly affects the
  - a. Hip
  - b. Knee
  - c. Elbow
  - d. Wrist
3. The most common neurologic finding of Lyme disease is
  - a. Cranial nerve palsies
  - b. Peripheral neuropathies
  - c. Aseptic meningitis
  - d. Radiculopathies
4. A finding on CSF analysis that is associated with persistent Lyme disease is
  - a. Increased glucose
  - b. Eosinopenia
  - c. Reduced protein
  - d. Mild pleocytosis
5. The preferred treatment for all patients except women who are pregnant or nursing and children younger than 12 years is oral
  - a. Amoxicillin
  - b. Doxycycline
  - c. Cefotaxime
  - d. Erythromycin
6. The Jarisch-Herxheimer-like reaction occurs most commonly with
  - a. Tetracycline
  - b. Clarithromycin
  - c. Ceftriaxone
  - d. Azithromycin

### NORMAL-PRESSURE HYDROCEPHALUS, PAGE 51

7. Normal-pressure hydrocephalus (NPH) is characterized by a clinical triad that includes urinary incontinence and
  - a. Seizures
  - b. Headache
  - c. Psychosis
  - d. Gait difficulty
8. The majority of cases of NPH are
  - a. Caused by a subarachnoid hemorrhage
  - b. Secondary to meningitis
  - c. Idiopathic
  - d. The result of a head injury
9. Cognitive symptoms of NPH are primarily subcortical features, such as
  - a. Aphasia
  - b. Depression
  - c. Delusions
  - d. Apathy
10. Multiple studies suggest a correlation between surgical outcome and
  - a. Duration of symptoms
  - b. Findings on lumbar puncture
  - c. Degree of cognitive impairment
  - d. Presence of dementia
11. The major immediate complication associated with shunt insertion is
  - a. Subdural fluid collection
  - b. Intracerebral hemorrhage
  - c. Infection
  - d. Seizures
12. The earliest and most likely response to shunt placement is improvement in
  - a. Incontinence
  - b. Cognition
  - c. Gait
  - d. Mood